

# DSA 8020 R Session 2: Multiple Linear Regression I

Whitney

## Contents

Species diversity on the Galapagos Islands . . . . .	1
First Step: Load the data . . . . .	1
Plot the pairwise scatterplots . . . . .	1
Correlation matrix . . . . .	2
Model 1: Fitting a simple linear regression . . . . .	3
Model 2: Adding <i>Area</i> . . . . .	4
Model 3: Adding <i>Adjacent</i> . . . . .	6
Full Model . . . . .	7
Parameter Estimation . . . . .	8
ANOVA . . . . .	9
Simulation . . . . .	9
$R^2$ vs. $R_{adj}^2$ . . . . .	9

## Species diversity on the Galapagos Islands


### First Step: Load the data

```
#installinstall.packages("faraway")
library(faraway)
data(gala)
head(gala)
```

```
##           Species Endemics  Area Elevation Nearest Scruz Adjacent
## Baltra           58      23 25.09      346      0.6  0.6      1.84
## Bartolome        31      21  1.24      109      0.6 26.3     572.33
## Caldwell          3       3  0.21      114      2.8 58.7      0.78
## Champion         25       9  0.10       46      1.9 47.4      0.18
## Coamano           2       1  0.05       77      1.9  1.9     903.82
## Daphne.Major     18      11  0.34      119      8.0  8.0      1.84
```

### Plot the pairwise scatterplots

```
pairs(gala[, -2], cex = 0.95, col = "blue", las = 1)
```



### Correlation matrix

```
cor(gala[, -2])
```

```
##           Species      Area  Elevation  Nearest      Scruz
## Species  1.0000000  0.6178431  0.73848666 -0.01409407 -0.17114244
## Area     0.61784307  1.0000000  0.75373492 -0.11110320 -0.10078493
## Elevation 0.73848666  0.7537349  1.00000000 -0.01107698 -0.01543829
## Nearest  -0.01409407 -0.1111032 -0.01107698  1.00000000  0.61541036
## Scruz    -0.17114244 -0.1007849 -0.01543829  0.61541036  1.00000000
## Adjacent  0.02616635  0.1800376  0.53645782 -0.11624788  0.05166066
##
##           Adjacent
## Species  0.02616635
## Area     0.18003759
## Elevation 0.53645782
## Nearest  -0.11624788
## Scruz    0.05166066
## Adjacent 1.00000000
```


## Model 1: Fitting a simple linear regression

Here we use *Elevation* as the predictor as it has the highest correlation with *Species*

```
M1 <- lm(Species ~ Elevation, data = gala)
summary(M1)
```

```
##
## Call:
## lm(formula = Species ~ Elevation, data = gala)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -218.319  -30.721  -14.690    4.634   259.180
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  11.33511   19.20529    0.590    0.56
## Elevation     0.20079    0.03465    5.795 3.18e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 78.66 on 28 degrees of freedom
## Multiple R-squared:  0.5454, Adjusted R-squared:  0.5291
## F-statistic: 33.59 on 1 and 28 DF,  p-value: 3.177e-06
```

```
plot(gala$Elevation, gala$Species, xlab = "Elevation (m)", ylab = "The number of plant species",
      las = 1, pch = 16, col = "blue")
abline(M1)
```



## Model 2: Adding Area

```
M2 <- lm(Species ~ Elevation + Area, data = gala)
summary(M2)
```


```
##
## Call:
## lm(formula = Species ~ Elevation + Area, data = gala)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -192.619  -33.534  -19.199    7.541  261.514
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  17.10519   20.94211    0.817  0.42120
## Elevation     0.17174    0.05317    3.230  0.00325 **
## Area           0.01880    0.02594    0.725  0.47478
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 79.34 on 27 degrees of freedom
## Multiple R-squared:  0.554, Adjusted R-squared:  0.521
## F-statistic: 16.77 on 2 and 27 DF,  p-value: 1.843e-05
```

```

library(fields)
Elevation_grid <- seq(0, 1750, 10)
Area_grid <- seq(0, 4700, 10)
temp <- expand.grid(Elevation_grid, Area_grid)
x_new <- data.frame(Elevation = temp$Var1, Area = temp$Var2)

y_pred <- matrix(predict(M2, x_new), nrow = length(Elevation_grid))
image.plot(Elevation_grid, Area_grid, y_pred, las = 1,
           xlab = "Elevation (m)", ylab = expression(paste("Area (", km^2, ")")))

```



```

library(plot3D)

```

```


## Warning in system2("/usr/bin/otool", c("-L", shQuote(DSO)), stdout = TRUE):
## running command ''/usr/bin/otool' -L '/Library/Frameworks/R.framework/Resources/
## library/tcltk/libs//tcltk.so'' had status 69

```

```

# fitted points for droplines to surface
fitpoints <- predict(M2)
# scatter plot with regression plane
scatter3D(gala$Elevation, gala$Elevation, gala$Species, pch = 16, cex = 1,
          theta = 20, phi = 30, ticktype = "detailed",
          xlab = "Elevation", ylab = "Area", zlab = "Species",
          surf = list(x = Elevation_grid, y = Area_grid, z = y_pred, facets = NA, fit = fitpoints))

```



### Model 3: Adding *Adjacent*

```
M3 <- lm(Species ~ Elevation + Area + Adjacent, data = gala)
summary(M3)
```

```
##
## Call:
## lm(formula = Species ~ Elevation + Area + Adjacent, data = gala)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -124.064  -34.283   -8.733   27.972  195.973
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  -5.71893   16.90706  -0.338  0.73789
## Elevation     0.31498    0.05211   6.044  2.2e-06 ***
## Area          -0.02031    0.02181  -0.931  0.36034
## Adjacent     -0.07528    0.01698  -4.434  0.00015 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Residual standard error: 61.01 on 26 degrees of freedom
## Multiple R-squared: 0.746, Adjusted R-squared: 0.7167
## F-statistic: 25.46 on 3 and 26 DF, p-value: 6.683e-08
```

## Full Model

```
M4 <- lm(Species ~ Elevation + Area + Adjacent + Nearest + Scruz, data = gala)
summary(M4)
```

```
##
## Call:
## lm(formula = Species ~ Elevation + Area + Adjacent + Nearest +
##     Scruz, data = gala)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -111.679  -34.898   -7.862   33.460  182.584
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  7.068221  19.154198   0.369 0.715351
## Elevation    0.319465   0.053663   5.953 3.82e-06 ***
## Area        -0.023938   0.022422  -1.068 0.296318
## Adjacent    -0.074805   0.017700  -4.226 0.000297 ***
## Nearest     0.009144   1.054136   0.009 0.993151
## Scruz       -0.240524   0.215402  -1.117 0.275208
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 60.98 on 24 degrees of freedom
## Multiple R-squared: 0.7658, Adjusted R-squared: 0.7171
## F-statistic: 15.7 on 5 and 24 DF, p-value: 6.838e-07
```

```
predict(M4)
```

```
##      Baltra      Bartolome      Caldwell      Champion      Coamano Daphne.Major
## 116.7259460 -7.2731544 29.3306594 10.3642660 -36.3839155 43.0877052
## Daphne.Minor      Darwin      Eden      Enderby      Espanola      Fernandina
## 33.9196678 -9.0189919 28.3142017 30.7859425 47.6564865 96.9895982
## Gardner1      Gardner2      Genovesa      Isabela      Marchena      Onslow
## -4.0332759 64.6337956 -0.4971756 386.4035578 88.6945404 4.0372328
## Pinta      Pinzon      Las.Plazas      Rabida SanCristobal SanSalvador
## 215.6794862 150.4753750 35.0758066 75.5531221 206.9518779 277.6763183
## SantaCruz      SantaFe      SantaMaria      Seymour      Tortuga      Wolf
## 261.4164131 85.3764857 195.6166286 49.8050946 52.9357316 26.7005735
```

```
confint(M4)
```

```
##              2.5 %      97.5 %
## (Intercept) -32.4641006 46.60054205
```

```
## Elevation      0.2087102  0.43021935
## Area          -0.0702158  0.02233912
## Adjacent      -0.1113362 -0.03827344
## Nearest       -2.1664857  2.18477363
## Scruz         -0.6850926  0.20404416
```

## Parameter Estimation

```
X <- model.matrix(M4)
y <- gala$Species
# regression parameters
(beta_hat <- solve(t(X) %*% X) %*% t(X) %*% y)
```

```
##                [,1]
## (Intercept)  7.068220709
## Elevation    0.319464761
## Area        -0.023938338
## Adjacent    -0.074804832
## Nearest     0.009143961
## Scruz       -0.240524230
```

```
beta_hat_faster <- solve(crossprod(X), crossprod(X, y))
# fitted values
(y_hat <- X %*% solve(t(X) %*% X) %*% t(X) %*% y)
```

```
##                [,1]
## Baltra       116.7259460
## Bartolome    -7.2731544
## Caldwell      29.3306594
## Champion     10.3642660
## Coamano      -36.3839155
## Daphne.Major  43.0877052
## Daphne.Minor  33.9196678
## Darwin       -9.0189919
## Eden         28.3142017
## Enderby      30.7859425
## Espanola     47.6564865
## Fernandina   96.9895982
## Gardner1     -4.0332759
## Gardner2     64.6337956
## Genovesa     -0.4971756
## Isabela     386.4035578
## Marchena     88.6945404
## Onslow       4.0372328
## Pinta       215.6794862
## Pinzon      150.4753750
## Las.Plazas   35.0758066
## Rabida       75.5531221
## SanCristobal 206.9518779
## SanSalvador  277.6763183
## SantaCruz    261.4164131
```



```
## SantaFe      85.3764857
## SantaMaria  195.6166286
## Seymour     49.8050946
## Tortuga     52.9357316
## Wolf        26.7005735
```

## ANOVA

```
anova(M4)
```

```
## Analysis of Variance Table
##
## Response: Species
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Elevation  1 207828  207828  55.8981 1.023e-07 ***
## Area       1   3307    3307   0.8895 0.3550197
## Adjacent   1  73171   73171  19.6804 0.0001742 ***
## Nearest    1   2909    2909   0.7823 0.3852165
## Scruz      1   4636    4636   1.2469 0.2752082
## Residuals 24  89231    3718
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```



## Simulation

$R^2$  vs.  $R_{adj}^2$

```
set.seed(123)
N = 500
x1 <- replicate(N, rnorm(30))
x2 <- replicate(N, rnorm(30))
y1 <- apply(x1, 2, function(x) 5 + 2 * x + rnorm(30, 0, 1))
R.sq <- array(dim = c(N, 4))
for (i in 1:N){
  R.sq[i, 1] = summary(lm(y1[, i] ~ x1[, i]))$r.squared
  R.sq[i, 2] = summary(lm(y1[, i] ~ x1[, i]))$adj.r.squared
  R.sq[i, 3] = summary(lm(y1[, i] ~ x1[, i] + x2[, i]))$r.squared
  R.sq[i, 4] = summary(lm(y1[, i] ~ x1[, i] + x2[, i]))$adj.r.squared
}

par(mfrow = c(1, 2))
plot(R.sq[, 1], R.sq[, 3], pch = 16, cex = 0.65, col = "blue",
     xlab = expression(paste("Model 1: ", R^2)),
     ylab = expression(paste("Model 2: ", R^2)))
abline(0, 1)

boxplot(R.sq[, 3] - R.sq[, 1], las = 1, xlab = expression(paste(R^2, " Model 2 - Model 1")))
abline(h = 0, lty = 2, col = "red")
```



```

par(las = 1, mfrow = c(1, 2), mar = c(5.1, 4.6, 1.1, 1.1))
plot(R.sq[, 2], R.sq[, 4], pch = 16, cex = 0.5, col = "blue",
     xlab = expression(paste("Model 1: ", R[adj]^2)),
     ylab = expression(paste("Model 2: ", R[adj]^2)))
abline(0, 1)

boxplot(R.sq[, 4] - R.sq[, 2], las = 1, xlab = expression(paste(R[adj]^2, " Model 2 - Model 1")))
abline(h = 0, lty = 2, col = "red")

```

