

Lecture 14

Regression with Time Series Errors, Unit Root Tests, Spurious Correlations, and Prewhitening

Reading: Cryer and Chen (2008): Chapter 3.3-3.4; Chapter 6.4; Chapter 11.3-11.4

MATH 4070: Regression and Time-Series Analysis

Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Whitney Huang
Clemson University

Agenda

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



- 1 Time Series Regression Models
- 2 Generalized Least Squares Regression
- 3 Unit Root Tests in Time Series Analysis
- 4 Spurious Correlation and Prewhitening

Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Suppose we have the following time series model for $\{Y_t\}$:

$$Y_t = m_t + \eta_t,$$

where

- m_t captures the mean of $\{Y_t\}$, i.e., $\mathbb{E}(Y_t) = m_t$
- $\{\eta_t\}$ is a zero mean stationary process with ACVF $\gamma_\eta(\cdot)$

The component $\{m_t\}$ may depend on time t , or possibly on other explanatory series

Example Models for m_t : Trends and Seasonality

- **Constant trend model:** For each t let $m_t = \beta_0$ for some unknown parameter β_0

- **Simple linear regression:** For unknown parameters β_0 and β_1 ,

$$m_t = \beta_0 + \beta_1 x_t,$$

where $\{x_t\}$ is some explanatory variable indexed in time (may just be a function of time or could be other series)

- **Harmonic regression:** For each t let

$$m_t = A \cos(2\pi\omega t + \phi),$$

where $A > 0$ is the amplitude (an unknown parameter), $\omega > 0$ is the frequency of the sinusoid (usually known), and $\phi \in (-\pi, \pi]$ is the phase (usually unknown). We can rewrite this model as

$$m_t = \beta_0 x_{1,t} + \beta_1 x_{2,t},$$

where $x_{1,t} = \cos(2\pi\omega t)$ and $x_{2,t} = \sin(2\pi\omega t)$

Multiple Linear Regression Model

Suppose there are p explanatory series $\{x_{j,t}\}_{j=1}^p$, the time series model for $\{Y_t\}$ is

$$Y_t = m_t + \eta_t,$$

where

$$m_t = \beta_0 + \sum_{j=1}^p \beta_j x_{j,t},$$

and $\{\eta_t\}$ is a mean zero stationary process with ACVF $\gamma_\eta(\cdot)$

We can write the linear model in matrix notation:

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta},$$

where $\mathbf{Y} = (Y_1, \dots, Y_n)^T$ is the observation vector, the coefficient vector is $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_p)^T$, $\boldsymbol{\eta} = (\eta_1, \dots, \eta_n)^T$ is the error vector, and the design matrix is

$$\mathbf{X} = \begin{bmatrix} 1 & x_{1,1} & x_{2,1} & \cdots & x_{p,1} \\ 1 & x_{1,2} & x_{2,2} & \cdots & x_{p,2} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 1 & x_{1,n} & x_{2,n} & \cdots & x_{p,n} \end{bmatrix}$$

Model Estimates & Distribution for i.i.d. Errors

Suppose $\{\eta_t\}$ is i.i.d. $N(0, \sigma^2)$. Then the ordinary least squares (OLS) estimate of β is

$$\hat{\beta}_{\text{OLS}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y},$$

with

$$\hat{\sigma}^2 = \frac{(\mathbf{Y} - \mathbf{X}\hat{\beta}_{\text{OLS}})^T (\mathbf{Y} - \mathbf{X}\hat{\beta}_{\text{OLS}})}{n - (p + 1)}$$



- Gauss-Markov theorem: $\hat{\beta}_{\text{OLS}}$ is the best linear unbiased estimator (BLUE) of β
- We have

$$\hat{\beta}_{\text{OLS}} \sim N(\beta, \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1})$$

is independent of

$$\frac{(n - (p + 1))\hat{\sigma}^2}{\sigma^2} \sim \chi_{n-(p+1)}^2$$

Temperatures and Tree Ring Proxies [Jones & Mann, 2004]



Residuals from a linear regression fit are **correlated in time** \Rightarrow
OLS is not appropriate here 😊

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Generalized Least Squares Regression

When dealing with time series the errors $\{\eta_t\}$ are typically correlated in time

- Assuming the errors $\{\eta_t\}$ are a stationary Gaussian process, consider the model

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta},$$

where $\boldsymbol{\eta}$ has a multivariate normal distribution, i.e.,
 $\boldsymbol{\eta} \sim N(\mathbf{0}, \Sigma)$

- The **generalized least squares (GLS) estimate** of $\boldsymbol{\beta}$ is

$$\hat{\boldsymbol{\beta}}_{\text{GLS}} = (\mathbf{X}^T \Sigma^{-1} \mathbf{X})^{-1} \mathbf{X}^T \Sigma^{-1} \mathbf{Y},$$

with

$$\hat{\sigma}^2 = \frac{(\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}_{\text{GLS}})^T (\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}_{\text{GLS}})}{n - (p + 1)}$$

Gauss-Markov theorem: β_{GLS} is the best linear unbiased estimator (BLUE) of β

- We have

$$\hat{\beta}_{\text{GLS}} \sim N(\beta, \sigma^2 (\mathbf{X}^T \Sigma^{-1} \mathbf{X})^T)$$

- The variance of linear combinations of $\hat{\beta}_{\text{GLS}}$ is less than or equal to the variance of linear combinations of $\hat{\beta}_{\text{OLS}}$, that is:

$$\text{Var}(\mathbf{c}^T \hat{\beta}_{\text{GLS}}) \leq \text{Var}(\mathbf{c}^T \hat{\beta}_{\text{OLS}})$$

The main problem in applying GLS in practice is that Σ depends on ϕ , θ , and σ^2 and we have to estimate these

- A two-step procedure
 - 1 Estimate β by OLS, calculating the residuals $\hat{\eta} = Y - X\hat{\beta}_{OLS}$, and fit an ARMA to $\hat{\eta}$ to get Σ
 - 2 Re-estimate β using GLS
- Alternatively, we can consider one-shot **maximum likelihood methods**

Model:

$$Y = X\beta + \eta,$$

where $\eta \sim N(\mathbf{0}, \Sigma)$

$$\Rightarrow Y \sim N(X\beta, \Sigma)$$

We maximize the **Gaussian** likelihood

$$\begin{aligned} L_n(\beta, \phi, \theta, \sigma^2) \\ = (2\pi)^{-n/2} |\Sigma|^{-1/2} \exp \left[-\frac{1}{2} (Y - X\beta)^T \Sigma^{-1} (Y - X\beta) \right] \end{aligned}$$

with respect to the regression parameters β and ARMA parameters ϕ, θ, σ^2 **simultaneously**

Comparison of Two-Step and One-Step Estimation Procedures

Let's conduct a Monte Carlo simulation with the following data-generating mechanism:

$$Y_t = 3 + 0.5x_t + \eta_t,$$

where $\eta_t = 0.8\eta_{t-1} + Z_t - 0.4Z_{t-1}$, $Z_t \sim N(0, 1)$.

- 1 Simulate 500 replications, each with 200 data points
- 2 Apply the two-step procedure: fit OLS, extract residuals, estimate ARMA model for $\hat{\Sigma}$, then refit using GLS.
- 3 Apply the one-step procedure to jointly estimate regression and ARMA parameters
- 4 Compare the estimation performance

Comparing Regression Slope Estimates

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening




Time Series
Regression Models

Generalized Least
Squares Regression


Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening



Method	OLS	GLS	MLE
Bias	-4e-4	9e-4	9e-4
Sd	0.046	0.035	0.035
CI coverage	90.8%	93.6%	93.6%
CI width	0.162	0.129	0.129

Comparing ARMA Estimates



Method	GLS	MLE
Bias	-0.038	-0.036
Sd	0.090	0.089
CI coverage	96.6%	96.2%
CI width	0.330	0.328

An Example: Lake Huron Levels

Model:

$$Y_t = m_t + \eta_t$$

where

$$m_t = \beta_0 + \beta_1 t$$

$\{\eta_t\}$ is some ARMA(p, q) process

- **Scientific Question:** Is there evidence that the lake level has changed linearly over the years 1875-1972?
- **Statistical Hypothesis:**

Fitting Result form the Two-Step Procedure

1 OLS:
`lm(formula = LakeHuron ~ years)`

Residuals:

Min	1Q	Median	3Q	Max
-2.50997	-0.72726	0.00083	0.74402	2.53565

Coefficients:

	Estimate	Std. Error	t value
(Intercept)	625.554918	7.764293	80.568
years	-0.024201	0.004036	-5.996

2 AR:
`arima(x = lm$residuals, order = c(2, 0, 0), include.mean = FALSE)`

Coefficients:

	ar1	ar2
	1.0050	-0.2925
s.e.	0.0976	0.1002

3 Refit GLS

Will leave it to you as an exercise

Fitting Result from One-Step MLE

```
> mle <- arima(LakeHuron, order = c(2, 0, 0),  
+             xreg = cbind(rep(1,length(LakeHuron)), years),  
+             include.mean = FALSE)  
> mle
```

Call:





```
arima(x = LakeHuron, order = c(2, 0, 0), xreg = cbind(rep(1, length(LakeHuron)),  
years), include.mean = FALSE)
```

Coefficients:

	ar1	ar2	rep(1, length(LakeHuron))	
	1.0048	-0.2913	620.5115	
s.e.	0.0976	0.1004	15.5771	
	years			
	-0.0216			
s.e.	0.0081			

sigma² estimated as 0.4566: log likelihood = -101.2, aic = 212.4

MLE Fit Diagnostics



```
> plot.residuals(years, resid(mle), xlab = "Year", ylab = "Residuals")
```

Box-Ljung test

data: y

X-squared = 6.2088, df = 19, p-value = 0.9974

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Comparing Confidence Intervals

Regression Slope β_1 :

Method	2.5%	Point Est.	97.5%
OLS	-0.0322	-0.0242	-0.0162
MLE	-0.0374	-0.0216	-0.0057



AR ϕ_1 :

Method	2.5%	Point Est.	97.5%
GLS	0.813	1.005	1.196
MLE	0.813	1.005	1.196

AR ϕ_2 :

Method	2.5%	Point Est.	97.5%
GLS	-0.489	-0.293	-0.096
MLE	-0.488	-0.291	-0.095

Unit Root Tests: Tests for Non-Stationarity

Suppose we have X_1, \dots, X_n that follow the model

$$(X_t - \mu) = \phi(X_{t-1} - \mu) + Z_t,$$

where $\{Z_t\}$ is a $WN(0, \sigma^2)$ process

- A **unit root test** considers the following hypotheses:

$$H_0 : \phi = 1 \text{ versus } H_a : |\phi| < 1$$

Unit Root Tests: Tests for Non-Stationarity

Suppose we have X_1, \dots, X_n that follow the model

$$(X_t - \mu) = \phi(X_{t-1} - \mu) + Z_t,$$

where $\{Z_t\}$ is a $WN(0, \sigma^2)$ process

- A **unit root test** considers the following hypotheses:

$$H_0 : \phi = 1 \text{ versus } H_a : |\phi| < 1$$

- Note that where $|\phi| < 1$ the process is **stationary** (and causal) while $\phi = 1$ leads to a nonstationary process

Unit Root Tests: Tests for Non-Stationarity

Suppose we have X_1, \dots, X_n that follow the model

$$(X_t - \mu) = \phi(X_{t-1} - \mu) + Z_t,$$

where $\{Z_t\}$ is a $WN(0, \sigma^2)$ process

- A **unit root test** considers the following hypotheses:

$$H_0 : \phi = 1 \text{ versus } H_a : |\phi| < 1$$

- Note that where $|\phi| < 1$ the process is **stationary** (and causal) while $\phi = 1$ leads to a nonstationary process
- **Exercise:** Letting $Y_t = \nabla X_t = X_t - X_{t-1}$, show that

$$\begin{aligned} Y_t &= (1 - \phi)\mu + (\phi - 1)X_{t-1} + Z_t \\ &= \phi_0^* + \phi_1^* X_{t-1} + Z_t, \end{aligned}$$

where $\phi_0^* = (1 - \phi)\mu$ and $\phi_1^* = (\phi - 1)$

Unit Root Tests via Ordinary Least Squares Argument

- We can estimate ϕ_0^* and ϕ_1^* using ordinary least squares

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Unit Root Tests via Ordinary Least Squares Argument

- We can estimate ϕ_0^* and ϕ_1^* using ordinary least squares
- Using the estimate of ϕ_1^* , $\hat{\phi}_1^*$, and its standard error, $\widehat{SE}(\hat{\phi}_1^*)$, the **Dickey-Fuller statistics** is

$$T = \frac{\hat{\phi}_1^*}{\widehat{SE}(\hat{\phi}_1^*)}$$

Unit Root Tests via Ordinary Least Squares Argument

- We can estimate ϕ_0^* and ϕ_1^* using ordinary least squares
- Using the estimate of ϕ_1^* , $\hat{\phi}_1^*$, and its standard error, $\widehat{SE}(\hat{\phi}_1^*)$, the **Dickey-Fuller statistics** is

$$T = \frac{\hat{\phi}_1^*}{\widehat{SE}(\hat{\phi}_1^*)}$$

- Under H_0 this statistic follows a **Dickey-Fuller distribution**. For a level α test we reject if the observed test statistic is smaller than a critical value C_α

α	0.01	0.05	0.10
C_α	-3.43	-2.86	-2.57

Unit Root Tests via Ordinary Least Squares Argument

- We can estimate ϕ_0^* and ϕ_1^* using ordinary least squares
- Using the estimate of ϕ_1^* , $\hat{\phi}_1^*$, and its standard error, $\widehat{SE}(\hat{\phi}_1^*)$, the **Dickey-Fuller statistics** is

$$T = \frac{\hat{\phi}_1^*}{\widehat{SE}(\hat{\phi}_1^*)}$$

- Under H_0 this statistic follows a **Dickey-Fuller distribution**. For a level α test we reject if the observed test statistic is smaller than a critical value C_α

α	0.01	0.05	0.10
C_α	-3.43	-2.86	-2.57

- We can extend to other processes (AR(p), ARMA(p, q), and MA(q))—see Brockwell and Davis [2016, Section 6.3] for further details



Unit Root Test: Simulated Examples

Recall

$$\nabla X_t = \phi_0^* + \phi_1^* X_{t-1} + Z_t,$$

where $\phi_0^* = (1 - \phi)\mu$ and $\phi_1^* = (\phi - 1)$

Let's demonstrate the test with a simulated **random walk** ($\phi = 1$)
and a simulated **white noise** ($\phi = 0$)



Unit Root Test: Simulated Examples Cont'd

```
> diff.rw <- diff(rw); n <- length(rw)
> ys <- diff.rw; xs <- rw[1:(n-1)]
> ols.rw <- lm(ys ~ xs); summary(ols.rw)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.10125	0.05973	1.695	0.0906 .
xs	-0.01438	0.00899	-1.600	0.1102

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Unit Root Test: Simulated Examples Cont'd

```
> diff.rw <- diff(rw); n <- length(rw)
> ys <- diff.rw; xs <- rw[1:(n-1)]
> ols.rw <- lm(ys ~ xs); summary(ols.rw)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.10125	0.05973	1.695	0.0906 .
xs	-0.01438	0.00899	-1.600	0.1102

```
> diff.wn <- diff(wn)
> ys <- diff.wn; xs <- wn[1:(n-1)]
> ols.wn <- lm(ys ~ xs); summary(ols.wn)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-0.001138	0.045329	-0.025	0.98
xs	-1.002420	0.044843	-22.354	<2e-16

Augmented Dickey-Fuller Test in R

Augmented Dickey-Fuller (ADF) Test: to check for the presence of a unit root in a time series and determine if the series is stationary

H_0 : The time series has a unit root (**non-stationary**)

H_1 : The time series is **stationary**

If p -value < significance level (e.g., 0.05), reject $H_0 \Rightarrow$ **stationary**

```
> library(tseries)
> adf.test(rw)
```

Augmented Dickey-Fuller Test

```
data: rw
Dickey-Fuller = -1.9203, Lag order = 7, p-value =
0.612
alternative hypothesis: stationary
```

```
> adf.test(wn)
Warning in adf.test(wn) : p-value smaller than printed
```

Augmented Dickey-Fuller Test

```
data: wn
Dickey-Fuller = -7.8953, Lag order = 7, p-value =
0.01
alternative hypothesis: stationary
```

Lagged Regression and Cross-Covariances

Consider the lagged regression model:

$$Y_t = \beta_0 + \beta_1 X_{t-d} + \varepsilon_t,$$

where X 's are iid random variables with variance σ_X^2 and the ε 's are also white noise with variance σ_ε^2 and are independent of the X 's

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Lagged Regression and Cross-Covariances

Consider the lagged regression model:

$$Y_t = \beta_0 + \beta_1 X_{t-d} + \varepsilon_t,$$

where X 's are iid random variables with variance σ_X^2 and the ε 's are also white noise with variance σ_ε^2 and are independent of the X 's

The **cross-covariance function** of $\{Y_t\}$ and $\{X_t\}$ is

$$\gamma_{XY}(h) = \mathbb{E}[(X_{t+h} - \mu_X)(Y_t - \mu_Y)],$$

and the **cross-correlation function (CCF)** is

$$\rho_{XY}(h) = \frac{\gamma_{XY}(h)}{\sqrt{\gamma_X(0)\gamma_Y(0)}}.$$

Lagged Regression and Cross-Covariances

Consider the lagged regression model:

$$Y_t = \beta_0 + \beta_1 X_{t-d} + \varepsilon_t,$$

where X 's are iid random variables with variance σ_X^2 and the ε 's are also white noise with variance σ_ε^2 and are independent of the X 's

The **cross-covariance function** of $\{Y_t\}$ and $\{X_t\}$ is

$$\gamma_{XY}(h) = \mathbb{E}[(X_{t+h} - \mu_X)(Y_t - \mu_Y)],$$

and the **cross-correlation function (CCF)** is

$$\rho_{XY}(h) = \frac{\gamma_{XY}(h)}{\sqrt{\gamma_X(0)\gamma_Y(0)}}.$$


If $d > 0$, we say X_t leads Y_t , and we have CCF is identically zero except for lag $h = -d$, where CCF is $\frac{\beta_1 \sigma_X}{\sqrt{\beta_1^2 \sigma_X^2 + \sigma_\varepsilon^2}}$

Lagged Regression and Its CCF

Consider the following regression model:

$$Y_t = X_{t-2} + \varepsilon_t,$$


where $X_t \stackrel{i.i.d}{\sim} N(0, 1)$, $\varepsilon_t \stackrel{i.i.d}{\sim} N(0, 0.25)$, and X 's and ε 's are independent to each other. The CCF is $\frac{1}{\sqrt{1+0.25}} = 0.8944$ when $h = -2$, and 0 otherwise



Spurious Correlations

- The lagged regression discussed earlier may be too restrictive, as X_t , Y_t , and ε_t could be temporally correlated


Example: X_t and Y_t are **independent**, but both follow an AR(1)



Spurious Correlations

- The lagged regression discussed earlier may be too restrictive, as X_t , Y_t , and ε_t could be temporally correlated
- Temporal dependence makes the horizon blue dashed lines ($\pm 1.96/\sqrt{n}$) unreliable


Example: X_t and Y_t are **independent**, but both follow an AR(1)



Spurious Correlations



- The lagged regression discussed earlier may be too restrictive, as X_t , Y_t , and ε_t could be temporally correlated
- Temporal dependence makes the horizon blue dashed lines ($\pm 1.96/\sqrt{n}$) unreliable
- This can lead to **spurious correlations**

Example: X_t and Y_t are **independent**, but both follow an AR(1)



Spurious Correlations: An Example with Milk and Electricity Data

Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



- **Observed Correlation:** Milk production and electricity usage show a high correlation due to shared seasonal patterns
- **Temporal Dependence:** Both series exhibit seasonality and autocorrelation, making raw correlations misleading
- **Key Takeaway:** Spurious correlations highlight the need for detrending and deseasonalizing in time series analysis



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening

Understanding Prewhitening

Prewhitening: A technique to remove autocorrelation in a time series before analyzing cross-correlations

Steps in Prewhitening:

- Fit a time series model (e.g., ARMA) to $\{X_t\}$ and filter it to obtain residuals

Understanding Prewhitening

Prewhitening: A technique to remove autocorrelation in a time series before analyzing cross-correlations

Steps in Prewhitening:

- Fit a time series model (e.g., ARMA) to $\{X_t\}$ and filter it to obtain residuals
- Apply the same model to $\{Y_t\}$ for consistent filtering

Understanding Prewhitening

Prewhitening: A technique to remove autocorrelation in a time series before analyzing cross-correlations

Steps in Prewhitening:

- Fit a time series model (e.g., ARMA) to $\{X_t\}$ and filter it to obtain residuals
- Apply the same model to $\{Y_t\}$ for consistent filtering
- Compute the cross-correlation of the residuals

Understanding Prewhitening

Prewhitening: A technique to remove autocorrelation in a time series before analyzing cross-correlations

Steps in Prewhitening:

- Fit a time series model (e.g., ARMA) to $\{X_t\}$ and filter it to obtain residuals
- Apply the same model to $\{Y_t\}$ for consistent filtering
- Compute the cross-correlation of the residuals

Understanding Prewhitening


Prewhitening: A technique to remove autocorrelation in a time series before analyzing cross-correlations

Steps in Prewhitening:

- Fit a time series model (e.g., ARMA) to $\{X_t\}$ and filter it to obtain residuals
- Apply the same model to $\{Y_t\}$ for consistent filtering
- Compute the cross-correlation of the residuals


```
x <- arima.sim(n = 100, list(ar = 0.9))
y <- arima.sim(n = 100, list(ar = 0.9))
par(las = 1, mgp = c(2.2, 1, 0), mar = c(3.6, 3.6, 0.8, 0.6), mfrow = c(1, 2))
ccf(x, y)
prewhiten(x, y)
...

```



Applying Prewhitening to the Milk and Electricity Data Example

```
> me.dif = ts.intersect(diff(diff(milk, 12)),  
+ diff(diff(log(electricity), 12)))  
> prewhiten(as.vector(me.dif[, 1]), as.vector(me.dif[, 2]), ylab = 'CCF')  
> par(las = 1, mgp = c(2.2, 1, 0), mar = c(3.6, 3.6, 0.8, 0.6))  
> prewhiten(as.vector(me.dif[, 1]), as.vector(me.dif[, 2]), ylab = 'CCF')
```



Regression with
Time Series Errors,
Unit Root Tests,
Spurious
Correlations, and
Prewhitening



Time Series
Regression Models

Generalized Least
Squares Regression

Unit Root Tests in
Time Series Analysis

Spurious Correlation
and Prewhitening