

Lecture 2

Estimating Trend and Seasonality

Readings: CC08 Chapter 3; SS17 Chapter 2; BD Chapter 1.5

MATH 8090 Time Series Analysis

Week 2

Whitney Huang
Clemson University

1 The Classical Decomposition Model

2 Trend Estimation

3 Estimating Seasonality

The Classical (Additive) Decomposition Model

- The additive model for a time series $\{Y_t\}$ is

$$Y_t = \mu_t + s_t + \eta_t,$$


where

- μ_t is the **trend** component
 - s_t is the **seasonal** component
 - η_t is the **random (noise)** component with $\mathbb{E}(\eta_t) = 0$
- Standard procedure:
 - (1) Estimate/remove the trend and seasonal components
 - (2) Analyze the remainder, the residuals $\hat{\eta}_t = y_t - \hat{\mu}_t - \hat{s}_t$
- We will focus on (1) for this week

Mauna Loa Atmospheric CO₂ Concentration Revisited

Monthly atmospheric concentrations of CO₂ at the Mauna Loa Observatory [Source: [Keeling & Whorf](#), Scripps Institution of Oceanography]

```
data{r}  
data(co2)  
par(mar = c(3.8, 4, 0.8, 0.6))  
plot(co2, las = 1, xlab = "", ylab = "")  
mtext("Time (year)", side = 1, line = 2)  
mtext(expression(paste("CO"[2], " Concentration (ppm)")), side = 2, line = 2.5)  
data{r}
```



Estimating Trend for Nonseasonal Model

- Assuming $s_t = 0$ (i.e., there is no “seasonal” variation), we have

$$Y_t = \mu_t + \eta_t,$$

with $\mathbb{E}(\eta_t) = 0$

- Methods for **estimating trends**
 - Least squares regression
 - Smoothing
- Alternatively, one can **remove trend** by **differencing time series**

- The additive nonseasonal time series model for $\{Y_t\}$ is

$$Y_t = \mu_t + \eta_t,$$

where the trend is assumed to be a linear combination of known covariate series $\{x_{it}\}_{i=1}^p$

$$\mu_t = \beta_0 + \sum_{i=1}^p \beta_i x_{it}.$$

- Here we want to **estimate** $\beta = (\beta_0, \beta_1, \dots, \beta_p)^T$ from the data $\{y_t, \{x_{it}\}_{i=1}^p\}_{t=1}^T$
- You're likely quite familiar with this formulation already \Rightarrow **Regression Analysis**

Some Examples of Covariate Series $\{x_{it}\}$

- **Simple linear regression model:**

$$\mu_t = \beta_0 + \beta_1 x_t,$$

for example, the temperature trend at time t could be a constant (β_0) plus a multiple (β_1) of the carbon dioxide level at time t (x_t)

- **Polynomial regression model:**

$$\mu_t = \beta_0 + \sum_{i=1}^p \beta_i t^i$$

- **Change point model:**

$$\mu_t = \begin{cases} \beta_0 & \text{if } t \leq t^*; \\ \beta_0 + \beta_1 & \text{if } t \geq t^*. \end{cases}$$

Parameter Estimation: Ordinary Least Squares

- Like in the linear regression setting, we can estimate the parameters via **ordinary least squares (OLS)**
- Specifically, we minimize the following objective function:

$$\ell_{ols} = \sum_{t=1}^T (y_t - \beta_0 - \sum_{k=1}^p x_{kt} \beta_k)^2.$$

- The estimates $\beta = (\beta_0, \beta_1, \dots, \beta_p)^T$ minimizing the above objective function are called the **OLS estimates of β** \Rightarrow they are easiest to express in **matrix form**

The Model and Parameter Estimates in Matrix Form

- Matrix representation:

$$Y = X\beta + \eta,$$

$$\text{where } Y = \begin{bmatrix} Y_1 \\ \vdots \\ Y_T \end{bmatrix}, X = \begin{bmatrix} 1 & x_{11} & x_{12} & \cdots & x_{1p} \\ 1 & x_{21} & x_{22} & \cdots & x_{2p} \\ 1 & \vdots & \cdots & \cdots & \vdots \\ 1 & x_{T1} & x_{T2} & \cdots & x_{Tp} \end{bmatrix}, \text{ and}$$


$$\eta = \begin{bmatrix} \eta_1 \\ \vdots \\ \eta_T \end{bmatrix}$$

- Assuming $X^T X$ is **invertible**, the OLS estimate of β can be shown to be

$$\hat{\beta} = (X^T X)^{-1} X^T Y,$$

and the `lm` function in R calculates OLS estimates

Lake Huron Example Revisited



Let's **assume** there is a **linear trend in time** \Rightarrow we need to estimate the **intercept** β_0 and **slope** β_1

Call:

```
lm(formula = LakeHuron ~ yr)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.50997	-0.72726	0.00083	0.74402	2.53565

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	625.554918	7.764293	80.568	< 2e-16 ***
yr	-0.024201	0.004036	-5.996	3.55e-08 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.13 on 96 degrees of freedom

Multiple R-squared: 0.2725, Adjusted R-squared: 0.2649


F-statistic: 35.95 on 1 and 96 DF, p-value: 3.545e-08

Plot the (Estimated) Trend $\hat{\mu}_t = \hat{\beta}_0 + \hat{\beta}_1 t$



$\hat{\beta}_1 = -0.0242$ (ft/yr) \Rightarrow there seems to be a decreasing trend

Plot the Residuals $\{\hat{\eta}_t = y_t - \hat{\beta}_0 - \hat{\beta}_1 t\}$




$\{\hat{\eta}_t\}$ seems to exhibit some temporal dependence structure, should we worry about the results we have (recall OLS makes an i.i.d. assumption)?

Statistical Properties of the OLS Estimates with Correlated Errors

- Assume the components of X are not random, the OLS estimates $\hat{\beta}$ are **unbiased** for β

Proof:


- Since $\{\eta_t\}$ is typically not an i.i.d. process (see the acf plot below), statistical inferences regarding β will be invalid



Smoothing or Local Averaging

In certain situations, we may want to relax the assumption on the trend \Rightarrow “non-parametric” approach

Here, we break the time series up into “small” blocks (each with 10 years of data) and average each block




Doing this gives a very rough estimate of the trend. **Can we do better?**

Moving Average Smoother

- A **moving average smoother** estimates the trend at time t by averaging the current observation and the q nearest observations from either side. That is

$$\hat{\mu}_t = \frac{1}{2q+1} \sum_{j=-q}^q y_{t-j}$$



- q is the “smoothing” parameter, which controls the smoothness of the estimated trend $\hat{\mu}_t$

- Let $\alpha \in [0, 1]$ be some fixed constant, defined


$$\hat{\mu}_t = \begin{cases} Y_1 & \text{if } t = 1; \\ \alpha Y_t + (1 - \alpha)\hat{\mu}_{t-1} & t = 2, \dots, T. \end{cases}$$

- For $t = 2, \dots, T$, we can rewrite $\hat{\mu}_t$ as

$$\sum_{j=0}^{t-2} \alpha(1 - \alpha)^j Y_{t-j} + (1 - \alpha)^{t-1} Y_1.$$

\Rightarrow it is a one-sided moving average filter with **exponentially decreasing weights**. One can alter α to control the amounts of smoothing (see next slide for an example)

α is the Smoothing Parameter for Exponential Smoothing



The smaller the α , the smoother the resulting trend

The final method we consider for removing trends is differencing

- We define the first order difference operator ∇ as

$$\nabla Y_t = Y_t - Y_{t-1} = (1 - B)Y_t,$$

where B is the **backshift operator** and is defined as $BY_t = Y_{t-1}$.

- Similarly the general order difference operator $\nabla^q Y_t$ is **defined recursively** as $\nabla[\nabla^{q-1} Y_t]$
- The backshift operator of power q is defined as $B^q Y_t = Y_{t-q}$

In next slide we will see an example regarding the relationship between ∇^q and B^q

The second order difference is given by

$$\nabla^2 Y_t = \nabla[\nabla Y_t]$$

The second order difference is given by

$$\begin{aligned}\nabla^2 Y_t &= \nabla[\nabla Y_t] \\ &= \nabla[Y_t - Y_{t-1}]\end{aligned}$$

The second order difference is given by

$$\begin{aligned}\nabla^2 Y_t &= \nabla[\nabla Y_t] \\ &= \nabla[Y_t - Y_{t-1}] \\ &= (Y_t - Y_{t-1}) - (Y_{t-1} - Y_{t-2})\end{aligned}$$

The second order difference is given by

$$\begin{aligned}\nabla^2 Y_t &= \nabla[\nabla Y_t] \\ &= \nabla[Y_t - Y_{t-1}] \\ &= (Y_t - Y_{t-1}) - (Y_{t-1} - Y_{t-2}) \\ &= Y_t - 2Y_{t-1} + Y_{t-2}\end{aligned}$$

The second order difference is given by

$$\begin{aligned}\nabla^2 Y_t &= \nabla[\nabla Y_t] \\ &= \nabla[Y_t - Y_{t-1}] \\ &= (Y_t - Y_{t-1}) - (Y_{t-1} - Y_{t-2}) \\ &= Y_t - 2Y_{t-1} + Y_{t-2} \\ &= (1 - 2B + B^2)Y_t\end{aligned}$$

In the next slide we will see an example of using differencing to
remove the trend

Removing Trend via Differencing

Consider a time series data with a linear trend (i.e., $\{Y_t = \beta_0 + \beta_1 t + \eta_t\}$) where η_t is a stationary time series. Then first order differencing results in a stationary series with no trend. To see why

$$\begin{aligned}\nabla Y_t &= Y_t - Y_{t-1} \\ &= (\beta_0 + \beta_1 t + \eta_t) - (\beta_0 + \beta_1(t-1) + \eta_{t-1}) \\ &= \beta_1 + \eta_t - \eta_{t-1}\end{aligned}$$

This is the sum of a stationary series and a constant, and therefore we have successfully remove the linear trend.

- A polynomial trend of order q can be removed by q -th order differencing
- By q -th order differencing a time series we are shortening its length by q
- Differencing does not allow you to estimate the trend, only to remove it. *Therefore it is not appropriate if the aim of the analysis is to describe the trend*

Seasonal Component Estimation

- Let's consider a situation where a time series consists of only a seasonal component (assuming the trend has been estimated/removed). In this scenario,

$$Y_t = s_t + \eta_t,$$

with $\{s_t\}$ having period d (i.e., $s_{t+jd} = s_t$ for all integers j and t), $\sum_{t=1}^d s_t = 0$ and $\mathbb{E}(\eta_t) = 0$

- Two methods to estimate $\{s_t\}$
 - Harmonic regression
 - Seasonal mean model
- A method to remove $\{s_t\} \Rightarrow$ Lag differencing

Harmonic Regression

- A harmonic regression model has the form

$$s_t = \sum_{j=1}^k A_j \cos(2\pi f_j t + \phi_j).$$


For each $j = 1, \dots, k$:

- $A_j > 0$ is the amplitude of the j -th cosine wave
 - f_j controls the frequency of the j -th cosine wave (how often waves repeats)
 - $\phi_j \in [-\pi, \pi]$ is the phase of the j -th wave (where it starts)
- The above can be expressed as

$$\sum_{j=1}^k (\beta_{1j} \cos(2\pi f_j t) + \beta_{2j} \sin(2\pi f_j t)),$$

where $\beta_{1j} = A_j \cos(\phi_j)$ and $\beta_{2j} = A_j \sin(\phi_j) \Rightarrow$ **if $\{f_j\}_{j=1}^k$ are known, we can use regression techniques to estimate the parameters $\{\beta_{1j}, \beta_{2j}\}_{j=1}^k$**

Monthly Average Temperature in Dubuque, IA [Cryer & Chan, 2008]



Let's assume that there is no trend in this time series. In this context, our goal is to estimate s_t , the seasonal component.


Use a Harmonic Regression to Model Annual Cycles

Model: $s_t = \beta_0 + \beta_1 \cos(2\pi t) + \beta_2 \sin(2\pi t)$

⇒ annual cycles can be modeled by a linear combination of **cos** and **sin** with 1-year period.

In R, we can easily create these harmonics using the `harmonic` function in the `TSA` package

```
harmonics <- harmonic(tempdub, 1)
```



```

```{r}
harReg <- lm(tempdub ~ harmonics)
summary(harReg)

```

Call:

```
lm(formula = tempdub ~ harmonics)
```

Residuals:

Min	1Q	Median	3Q	Max
-11.1580	-2.2756	-0.1457	2.3754	11.2671


Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	46.2660	0.3088	149.816	< 2e-16	***
harmonicscos(2*pi*t)	-26.7079	0.4367	-61.154	< 2e-16	***
harmonicssin(2*pi*t)	-2.1697	0.4367	-4.968	1.93e-06	***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

# The Harmonic Regression Model Fit





- **Harmonics regression** assumes the seasonal pattern has a regular shape, i.e., the height of the peaks is the same as the depth of the troughs
- A less restrictive approach is to model  $\{s_t\}$  as

$$s_t = \begin{cases} \beta_1 & \text{for } t = 1, 1 + d, 1 + 2d, \dots & ; \\ \beta_2 & \text{for } t = 2, 2 + d, 2 + 2d, \dots & ; \\ \vdots & \vdots & ; \\ \beta_d & \text{for } t = d, 2d, 3d, \dots & . \end{cases}$$

- This is the **seasonal means** model, the parameters  $(\beta_1, \beta_2, \dots, \beta_d)^T$  can be estimated under the linear model framework (think about ANOVA)

## R Output

```
Call:
lm(formula = tempdub ~ month - 1)
```

Residuals:


Min	1Q	Median	3Q	Max
-8.2750	-2.2479	0.1125	1.8896	9.8250

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
monthJanuary	16.608	0.987	16.83	<2e-16	***
monthFebruary	20.650	0.987	20.92	<2e-16	***
monthMarch	32.475	0.987	32.90	<2e-16	***
monthApril	46.525	0.987	47.14	<2e-16	***
monthMay	58.092	0.987	58.86	<2e-16	***
monthJune	67.500	0.987	68.39	<2e-16	***
monthJuly	71.717	0.987	72.66	<2e-16	***
monthAugust	69.333	0.987	70.25	<2e-16	***
monthSeptember	61.025	0.987	61.83	<2e-16	***
monthOctober	50.975	0.987	51.65	<2e-16	***
monthNovember	36.650	0.987	37.13	<2e-16	***
monthDecember	23.642	0.987	23.95	<2e-16	***

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

# The Seasonal Means Model Fit



- The lag- $d$  difference operator,  $\nabla_d$ , is defined by


$$\nabla_d Y_t = Y_t - Y_{t-d} = (1 - B^d)Y_t.$$

**Note:** This is NOT  $\nabla^d$ !

- **Example:** Consider data that arise from the model  $Y_t = \beta_0 + \beta_1 t + s_t + \eta_t$ , which has a linear trend and seasonal component that repeats itself every  $d$  time points. Then by just seasonal differencing (lag- $d$  differencing here) this series becomes stationary.

$$\begin{aligned}\nabla_d Y_t &= Y_t - Y_{t-d} \\ &= [\beta_0 + \beta_1 t + s_t + \eta_t] - [\beta_0 + \beta_1(t-d) + s_{t-d} + \eta_{t-d}] \\ &= d\beta_1 + \eta_t - \eta_{t-d}\end{aligned}$$


# Estimating the Trend and Seasonal variation Together



Let's perform a regression analysis to model both  $\mu_t$  (assuming a linear time trend) and  $s_t$  (using `cos` and `sin`)

```
```\r\ntime <- as.numeric(time(co2))\nharmonics <- harmonic(co2, 1)\n\nlm_trendSeason <- lm(co2 ~ time + harmonics)\nsummary(lm_trendSeason)
```

The Regression Fit



Seasonal and Trend decomposition using Loess [Cleveland, et. al., 1990]

```
```{r}  
Seasonal and Trend decomposition using Loess (STL)
par(mar = c(4, 3.6, 0.8, 0.6))
stl <- stl(co2, s.window = "periodic")
plot(stl, las = 1)
```
```

