

MATH 8090: Stationary processes and Linear Processes

Whitney Huang, Clemson University

9/12-9/14/2023

Contents

Estimation of the Mean and Autocovariance Function of Stationary Processes	1
Mean Estimation	1
Autocovariance Function (ACVF)	2
Autocorrelation Function (ACF)	3
Lake Huron Example	4
Box test for temporal independence	7
Box and Pierce test Box and Pierce (1970)	7
Ljung-Box Test	7
Examples of MA(q) processes	8
Examples of AR(2) processes	10
References	12

Estimation of the Mean and Autocovariance Function of Stationary Processes

Mean Estimation

Given a stationary process $\{\eta_t\}_{t=1}^T$, the point estimator is $\bar{\eta} = \frac{1}{T} \sum_{t=1}^T \eta_t$. The variance of this estimator is

$$\nu_T = \text{Var}(\bar{\eta}) = \text{Var}\left(\frac{1}{T} \sum_{t=1}^T \eta_t\right) = \frac{1}{T} \sum_{h=-(T-1)}^{T-1} \left(1 - \frac{|h|}{T}\right) \gamma(h)$$


```
# Monte Carlo approximation
M = 1000; T = 200; phi = 0.9
set.seed(123)
sim <- replicate(M, arima.sim(n = T, list(ar = c(phi))))
eta_bar <- apply(sim, 2, mean)
hist(eta_bar, 40, col = "lightblue", border = "gray", las = 1, prob = T, main = "")
(nu_T_hat <- var(eta_bar))

## [1] 0.4659671
```

```

# Theoretical sampling dist
mu = 0
h <- -(T-1):(T-1)
nu_T <- (1 / T) * sum((1 - (abs(h) / T)) * (phi^(abs(h)) / (1 - phi^2)))
## Superimpose the true density curve
xg <- seq(-2.5, 2.5, 0.01)
abline(v = mu, lty = 2)
lines(xg, dnorm(xg, sd = sqrt(nu_T)), col = "red")

```



```

## Compare nu_T and nu
(nu_T <- (1 / T) * sum((1 - (abs(h) / T)) * (phi^(abs(h)) / (1 - phi^2))))

```

```

## [1] 0.4763158

```

```

(nu <- (1 / T) * (1 / (1 - phi)^2))

```

```

## [1] 0.5

```

Autocovariance Function (ACVF)

```
gamma <- apply(sim, 2, function(x) acf(x, plot = F, type = "covariance")$acf)
```

```
(est <- apply(gamma, 1, mean))
```

```


## [1] 4.762396047 4.210822098 3.718189717 3.276955489 2.883257698 2.529033768
## [7] 2.215391725 1.934657288 1.684410252 1.460895692 1.261443145 1.085257775
## [13] 0.927192703 0.791189571 0.671906177 0.563884323 0.468728429 0.380609541
## [19] 0.300366800 0.227258194 0.161799558 0.106047046 0.056009807 0.007812902

```

```
(true <- ARMAacf(ar = c(0.9), lag.max = 23) * (1 / (1 - phi^2)))
```

```
##          0           1           2           3           4           5           6           7
## 5.2631579 4.7368421 4.2631579 3.8368421 3.4531579 3.1078421 2.7970579 2.5173521
##          8           9          10          11          12          13          14          15
## 2.2656169 2.0390552 1.8351497 1.6516347 1.4864712 1.3378241 1.2040417 1.0836375
##         16          17          18          19          20          21          22          23
## 0.9752738 0.8777464 0.7899718 0.7109746 0.6398771 0.5758894 0.5183005 0.4664704
```

```
# The sample ACVF is a biased estimator
rg <- range(true, est)
plot(true, est, pch = 16, xlim = rg, ylim = rg, las = 1)
abline(0, 1, col = "blue")
```



```
# ACVF is non-negative definite
```

```
h <- outer(1:24, 1:24, "-")
Sigma <- phi^abs(h) / (1 - phi^2)
eigen(Sigma)$values
```

```
## [1] 64.7315984 26.5011371 11.7018213 6.1924112 3.7670568 2.5262589
## [7] 1.8165664 1.3759195 1.0849122 0.8833909 0.7385536 0.6313427
## [13] 0.5500994 0.4873821 0.4382698 0.3994115 0.3684691 0.3437787
## [19] 0.3241390 0.3086742 0.2967447 0.2878873 0.2817747 0.2781898
```

Autocorrelation Function (ACF)

Population ACF

$$\rho(h) = \text{Cor}(\eta_t, \eta_{t+h}) = \frac{\mathbb{E}[(\eta_t - \mu)(\eta_{t+h} - \mu)]}{\sqrt{\text{Var}(\eta_t)\text{Var}(\eta_{t+h})}}$$


Sample ACF

$$\hat{\rho}(h) = \frac{\hat{\gamma}(h)}{\hat{\gamma}(0)},$$

where $\hat{\gamma}(h) = \frac{1}{T} \sum_{t=1}^{T-|h|} (\eta_t - \bar{\eta})(\eta_{t+h} - \bar{\eta})$.


```
AR1 <- arima.sim(n = 100, list(ar = c(0.75)))
par(las = 1)
acf(AR1, lag = 100)
acf_true <- ARMAacf(ar = c(0.75), lag.max = 100)
points(0:100, acf_true, pch = 16, cex = 0.5, col = "red")
```

Series AR1




Lake Huron Example

```
data(LakeHuron)
par(las = 1, mfrow = c(2, 1), mar = c(4, 4, 0.8, 0.6))
plot(LakeHuron, ylab = "Depth (ft)", xlab = "Year")
points(LakeHuron, cex = 0.8, col = "blue", pch = 16)
acf(LakeHuron)
```




```
# Let's remove the (linear trend)
yr <- 1875:1972
lm <- lm(LakeHuron ~ yr)
plot(lm$residuals, ylab = "Depth (ft)", xlab = "Year", type = "l")
points(lm$residuals, cex = 0.8, col = "blue", pch = 16)
acf(lm$residuals)
```



```

par(mfrow = c(1, 1), las = 1)
plot(0:19, acf(LakeHuron, plot = F)$acf, type = "h", xlab = "Lag", ylab = "ACF", ylim = c(-0.2, 1))
abline(h = 0, col = "gray")
abline(h = c(-1, 1) * qnorm(0.975) / sqrt(length(LakeHuron)) , col = "blue", lty = 2)
acf_detrend <- acf(lm$residuals, plot = F)$acf
points(0:19, acf_detrend, col = "red")

```



Box test for temporal independence

Box and Pierce test Box and Pierce (1970)

We wish to test:

$$H_0 : \{\eta_1, \eta_2, \dots, \eta_T\} \text{ is an i.i.d. noise sequence}$$

$$H_1 : H_0 \text{ is false}$$

- Under H_0 ,

$$\hat{\rho}(h) \stackrel{\sim}{\sim} N(0, \frac{1}{T}) \stackrel{d}{=} \frac{1}{\sqrt{n}} N(0, 1)$$

- Hence

$$Q = T \sum_{i=1}^k \hat{\rho}^2(h) \stackrel{\sim}{\sim} \chi_{df=k}^2$$

- We reject H_0 if $Q > \chi_k^2(1 - \alpha)$, the $1 - \alpha$ quantile of the chi-squared distribution with k degrees of freedom

Ljung-Box Test

$$Q_{LB} = T(T-2) \sum_{h=1}^k \frac{\hat{\rho}^2(h)}{n-h} \stackrel{\sim}{\sim} \chi_k^2.$$


The Ljung-Box test Ljung and Box (1978) can be more powerful than the Box and Pierce test

```

out <- numeric(10000)
for (i in 1:10000) out[i] <- Box.test(rnorm(100), 4)$p.value
hist(out, 50, prob = T)

```

Histogram of out



```

out <- Box.test(rnorm(100), 4)
Box.test(lm$residuals, 4)

```

```

##
##  Box-Pierce test
##
## data: lm$residuals
## X-squared = 86.581, df = 4, p-value < 2.2e-16

```

```
Box.test(LakeHuron, 4, type = "Ljung")
```

```

##
##  Box-Ljung test
##
## data: LakeHuron
## X-squared = 143.87, df = 4, p-value < 2.2e-16


```

Examples of MA(q) processes

```


ma3 <- arima.sim(n = 200, list(ma = c(0.6, 0.5, 0.2)))
par(las = 1)
ts.plot(ma3)

```




```
acf(ma3)
acf_true <- ARMAacf(ma = c(0.6, 0.5, 0.2), lag.max = 20)
points(0:20, acf_true, col = "red")
```

Series ma3




Examples of AR(2) processes

```
ar2_1 <- arima.sim(n = 200, list(ar = c(0.5, 0.25)))
ts.plot(ar2_1)
```




```
acf(ar2_1)
points(0:20, ARMAacf(ar = c(0.5, 0.25), lag.max = 20), col = "red")
```

Series ar2_1




```
ar2_2 <- arima.sim(n = 200, list(ar = c(1.5, -0.75)))
ts.plot(ar2_2)
```



```
acf(ar2_2)
points(0:20, ARMAacf(ar = c(1.5, -0.75), lag.max = 20), col = "red")
```

Series ar2_2



References

- Box, George EP, and David A Pierce. 1970. "Distribution of Residual Autocorrelations in Autoregressive-Integrated Moving Average Time Series Models." *Journal of the American Statistical Association* 65 (332): 1509–26.
- Ljung, Greta M, and George EP Box. 1978. "On a Measure of Lack of Fit in Time Series Models." *Biometrika* 65 (2): 297–303.